

## Improving Acute Cardiac Syndrome Outcomes by Adopting the Occlusion Myocardial Infarction (OMI/NOMI) Classification

### Melhorando os Desfechos da Síndrome Coronária Aguda com a Integração da Classificação OMI (OCA/NOCA)

Bernardo Vidal Pimentel<sup>1</sup> , Marta Azevedo Ferreira<sup>2</sup> , José Nunes de Alencar<sup>3</sup> 

**Keywords:** Acute Coronary Syndrome; Non-ST Elevated Myocardial Infarction; ST Elevation Myocardial Infarction.

**Palavras-chave:** Infarte do Miocárdio com Supradesnível do Segmento ST; Infarte do Miocárdio sem Supradesnível do Segmento ST; Síndrome Coronária Aguda

We read with great interest the clinical case report by Rocha *et al*, where the diagnosis of an acute coronary occlusion (ACO) was suspected in a patient with an acute cardiac syndrome (ACS) without classic ST-segment elevation through the identification of the Aslanger Pattern, a pattern suggestive of an inferior ACO.<sup>1</sup> We congratulate the authors on their astute recognition and commend the team's effective management of the patient.

We would like to take this opportunity to discuss some numerical data presented, to contribute to a deeper understanding of the occlusion myocardial infarction (OMI) paradigm. Specifically, we believe that some of the figures regarding the diagnostic accuracy of electrocardiograms (ECGs) for ACO might be even more concerning than reported. The authors mention that 15% to 35% of patients had false-positive ECGs for STEMI, citing Meyers<sup>2</sup> and Aslanger (the DIFOCULT study).<sup>3</sup> However, this range may not fully reflect the current evidence. In a recent meta-analysis,<sup>4</sup> which included these two studies, there was found a pooled specificity of 96.5% for ECG criteria in diagnosing OMI. This translates to a false-positive rate of only 3.5%, a significantly lower figure than the 15% to 35% reported. More importantly, there was observed a low pooled sensitivity of 43.6%, indicating that more than half of the patients presenting with ACO do not exhibit ST-segment elevation. When we evaluated additional signs indicative of ACO, were evaluated, sensitivity increased to

78.1%, while specificity remained high at 94.4%. The positive likelihood ratio improved from 12.51 to 13.85. These findings suggest that ECGs, when interpreted with comprehensive OMI-specific criteria, are more reliable than previously thought for the early detection of OMI.

The timely recognition of patients with ACS remains a challenge for acute care physicians such as internal medicine doctors, and the misdiagnosis of acute myocardial infarction (AMI) can result in significant morbimortality. While the ECG is an essential diagnostic tool in the evaluation of patients with chest pain, relying solely on ST-segment elevation criteria may not be sufficient.

To overcome this identification problem, "OMI Manifesto" provided a theoretical framework for the early recognition and management of AMI.<sup>5</sup> This classification (Fig. 1) encompasses both STEMI and STEMI equivalents, aiming for a quicker identification to shorten door-to-balloon time and improve patient outcomes.<sup>6</sup> Additionally, the aforementioned observational studies found a beneficial advantage of the OMI/non-OMI classification compared to the traditional STEMI/non-STEMI.<sup>2,3</sup> Recognizing the importance of this paradigm, some of us collaborated with the original proponents to launch a Portuguese version of the OMI Manifesto.<sup>7</sup> In this version, the terms "Oclusão Coronária Aguda (OCA)" and "Ausência de Oclusão Coronária Aguda (NOCA)" were adopted to better reflect the concept in our language and facilitate its understanding within the Portuguese-speaking medical community.


Therefore, we believe that this case report published in *Revista da Sociedade Portuguesa de Medicina Interna*, holds significant importance in alerting the Portuguese medical community to the necessity of a paradigm shift from the traditional STEMI/non-STEMI classification to OMI/non-OMI – or OCA/NOCA in Portuguese – such as it has been alerted in the international community.<sup>8</sup> Such an approach is in alignment with the recent expert consensus statement from the American College of Cardiology, which urges clinicians dealing with acute chest pain not to overlook "STEMI equivalents,"<sup>9</sup> as well as with the latest guidelines from the European Society of Cardiology that identify non-STEMI patients with high-risk criteria potentially representing acute occlusion.<sup>10</sup> However, more concerted efforts from these

<sup>1</sup> Internal Medicine Department, Hospital da Luz Lisboa, Lisbon, Portugal

<sup>2</sup> Internal Medicine Department, Hospital Beatriz Ângelo, Lisbon, Portugal

<sup>3</sup> Electrocardiography Unit, Instituto Dante Pazzanese de Cardiologia, São Paulo, Brazil

<https://doi.org/10.24950/rspm.2649>



ACS – Acute coronary syndrome | ECG – Electrocardiogram | MI – Myocardial infarction | OMI – Occlusion myocardial infarction | STEMI – ST elevation myocardial infarction

**Figure 1:** Visual theoretical conception of OMI/NOMI (or OCA/NOCA) and STEMI/NSTEMI (not a diagnostic algorithm).

institutions are still needed to fully embrace and promote this paradigm shift. By redefining how we classify and name this condition, we can improve our diagnostic capabilities and, consequently, our treatment strategies, ultimately leading to better patient outcomes. ■

### Contributorship Statement

BVP - Original idea, conception, writing and creation of the figure and revision

MAF, JNA - Conception, writing and revision

All authors approved the final version to be published.

### Declaração de Contribuição

BVP – Ideia original, concepção, escrita e criação da figura e revisão

MAF, JNA - Concepção, escrita e revisão

Todos os autores aprovaram a versão final a ser publicada.

### Ethical Disclosures

Conflicts of Interest: The authors have no conflicts of interest to declare.

Financing Support: This work has not received any contribution, grant or scholarship

Confidentiality of Data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Protection of Human and Animal Subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki as

revised in 2024).

Provenance and Peer Review: Not commissioned; externally peer reviewed.

### Responsabilidades Éticas

Conflitos de Interesse: Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho.

Fontes de Financiamento: Não existiram fontes externas de financiamento para a realização deste artigo.

Confidencialidade dos Dados: Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes.

Proteção de Pessoas e Animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pela Comissão de Ética responsável e de acordo com a Declaração de Helsinquia revista em 2024 e da Associação Médica Mundial.

Proveniência e Revisão por Pares: Não comissionado; revisão externa por pares.

© Autor (es) (ou seu (s) empregador (es)) e Revista SPMI 2025. Reutilização permitida de acordo com CC BY-NC 4.0. Nenhuma reutilização comercial.

© Author(s) (or their employer(s)) and SPMI Journal 2025. Re-use permitted under CC BY-NC 4.0. No commercial re-use.

### Corresponding author / Autor correspondente:

Bernardo Vidal Pimentel - bernardo\_pimentel@hotmail.com

Internal Medicine Department, Hospital da Luz Lisboa, Lisbon, Portugal

Av. Lusíada 100, 1500-6500 Lisboa

Recebido / Received: 2024/11/07

Aceite / Accepted: 2025/01/15

Publicado / Published: ahead of print: 2025/00/00

## REFERENCES

1. Rocha D, Gouveia J, Ferreira C, Franco F "Acute Inferior Myocardial Infarction: Beyond the ST-Segment Elevation." *SPMI Case Rep.* 2024; 36-9. doi: [org/10.60591/crspmi.173](https://doi.org/10.60591/crspmi.173)
2. Pendell Meyers H, Bracey A, Lee D, Lichtenheld A, Li WJ, Singer DD, et al. Accuracy of OMI ECG findings versus STEMI criteria for diagnosis of acute coronary occlusion myocardial infarction. *Int J Cardiol Heart Vasc.* 2021;33:100767. doi: [10.1016/j.ijcha.2021.100767](https://doi.org/10.1016/j.ijcha.2021.100767).
3. Aslanger EK, Yıldırım Türk Ö, Simsek B, Bozbeyoglu E, Simsek MA, Yücel Karabay C, et al. Diagnostic accuracy of electrocardiogram for acute coronary Occlusion resulting in myocardial infarction (DIFOCULT Study). *Int J Cardiol Heart Vasc.* 2020;30:100603. doi: [10.1016/j.ijcha.2020.100603](https://doi.org/10.1016/j.ijcha.2020.100603).
4. de Alencar Neto JN, Scheffer MK, Correia BP, Franchini KG, Felicioni SP, De Marchi MF. Systematic review and meta-analysis of diagnostic test accuracy of ST-segment elevation for acute coronary occlusion. *Int J Cardiol.* 2024;402:131889. doi: [10.1016/j.ijcard.2024.131889](https://doi.org/10.1016/j.ijcard.2024.131889).
5. Meyers HP, Weingart SD, Smith SW. The OMI manifesto. *Dr. Smith's ECG Blog.* [accessed: May 18, 2023] Available at: <http://hqmeded-ecg.blogspot.com/2018/04/the-omi-manifesto.html>.
6. McLaren JTT, Meyers HP, Smith SW, Chartier LB. From STEMI to occlusion MI: paradigm shift and ED quality improvement. *CJEM.* 2022;24:250-5. doi: [10.1007/s43678-021-00255-z](https://doi.org/10.1007/s43678-021-00255-z).
7. Alencar JN, Feres F, Marchi MF, Franchini KG, Scheffer MK, Felicioni SP, et al. Beyond STEMI-NSTEMI Paradigm: Dante Pazzanese's Proposal for Occlusion Myocardial Infarction Diagnosis. *Arq Bras Cardiol.* 2024;121:e20230733. doi: [10.36660/abc.20230733](https://doi.org/10.36660/abc.20230733).
8. McLaren J, de Alencar JN, Aslanger EK, Meyers HP, Smith SW. From ST-Segment Elevation MI to Occlusion MI: The New Paradigm Shift in Acute Myocardial Infarction. *JACC Adv.* 2024;3:101314. doi: [10.1016/j.jaccadv.2024.101314](https://doi.org/10.1016/j.jaccadv.2024.101314).
9. Kontos MC, de Lemos JA, Deitelzweig SB, Diercks DB, Gore MO, Hess EP, et al. 2022 ACC Expert Consensus Decision Pathway on the Evaluation and Disposition of Acute Chest Pain in the Emergency Department: A Report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol.* 2022;80:1925-60. doi: [10.1016/j.jacc.2022.08.750](https://doi.org/10.1016/j.jacc.2022.08.750). Erratum in: *J Am Coll Cardiol.* 2024;84:771. doi: [10.1016/j.jacc.2024.07.009](https://doi.org/10.1016/j.jacc.2024.07.009).
10. Byrne RA, Rossello X, Coughlan JJ, Barbato E, Berry C, Chieffo A, et al. 2023 ESC Guidelines for the management of acute coronary syndromes. *Eur Heart J Acute Cardiovasc Care.* 2024;13:55-161. doi: [10.1093/ehjacc/zuad107](https://doi.org/10.1093/ehjacc/zuad107). Erratum in: *Eur Heart J Acute Cardiovasc Care.* 2024;13:455. doi: [10.1093/ehjacc/zuad156](https://doi.org/10.1093/ehjacc/zuad156).